Mon. Not. R. Astron. Soc. **000**, 1–6 (2011) Printed 20 July 2011 (MN LaTeX style file v2.2)

arXiv:1107.3770v1 [astro-ph.HE] 19 Jul 2011

# *Swift* monitoring of the central X-ray source in RCW 103

P. Esposito,[1]⋆ R. Turolla,[2,3] A. De Luca,[4,5] G. L. Israel,[6] A. Possenti[1] and D. N. Burrows[7]
[1] *INAF – Osservatorio Astronomico di Cagliari, località Poggio dei Pini, strada 54, I-09012 Capoterra, Italy*
[2] *Università di Padova, Dipartimento di Fisica, via F. Marzolo 8, I-35131 Padova, Italy*
[3] *Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK*
[4] *IUSS – Istituto Universitario di Studi Superiori, viale Lungo Ticino Sforza 56, I-27100 Pavia, Italy*
[5] *INAF – Istituto di Astrofisica Spaziale e Fisica Cosmica - Milano, via E. Bassini 15, I-20133 Milano, Italy*
[6] *INAF – Osservatorio Astronomico di Roma, via Frascati 33, I-00040 Monteporzio Catone, Italy*
[7] *Department of Astronomy & Astrophysics, The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802, USA*

Accepted 2011 July 19. Received 2011 July 19; in original form 2011 July 5

## ABSTRACT

The X-ray source 1E 161348–5055 lies at the centre of the 2-kyr-old supernova remnant RCW 103. Owing to its 24-ks modulation, orders-of-magnitude flux variability over a few months/years, and lack of an obvious optical counterpart, 1E 161348–5055 defies assignment to any known class of X-ray sources. Starting from April 2006, *Swift* observed 1E 161348–5055 with its X-ray telescope for ∼2 ks approximately once per month. During the five years covered, the source has remained in a quiescent state, with an average observed flux of $\sim 1.7 \times 10^{-12}$ erg cm$^{-2}$ s$^{-1}$ (1–10 keV), ∼20 times lower than the historical maximum attained in its 1999–2000 outburst. The long time-span of the *Swift* data allows us to obtain an accurate measure of the period of 1E 161348–5055 [$P = 24\,030.42(2)$ s] and to derive the first upper limit on its period derivative ($|\dot{P}| < 1.6 \times 10^{-9}$ s s$^{-1}$ at 3$\sigma$).

**Key words:** pulsars: general – stars: neutron – X-rays: individual: 1E 161348–5055.

## 1 INTRODUCTION

1E 161348–5055 (hereafter 1E 1613) was discovered with the *Einstein* satellite (Tuohy & Garmire 1980) close to the geometrical centre of the young supernova remnant (SNR) RCW 103 (age ∼2 kyr; Carter, Dickel & Bomans 1997). It was proposed as the first example of a radio-quiet (possibly owing to an unfavourable radio beaming), isolated, cooling neutron star (Tuohy & Garmire 1980; Tuohy et al. 1983; Gotthelf, Petre & Hwang 1997).

At present, there is little doubt that 1E 1613 indeed is a neutron star (De Luca et al. 2006) and the source is traditionally included in the class of the 'central compact objects' (CCOs; see De Luca 2008 for a review). CCOs are a small group of young and seemingly isolated X-ray-emitting neutron stars (with thermal-like spectra), observed close to the centre of non-plerionic SNRs and without obvious counterparts in other wavebands. However, its peculiar temporal behaviour distinguishes 1E 1613 from the other CCOs (actually, it singles this source out as a unique object in general). The first peculiarity of the source is its orders-of-magnitude X-ray flux variability on a few months/years time-scale (Gotthelf, Petre & Vasisht 1999; Garmire et al. 2000; Sanwal et al. 2002; Becker & Aschenbach 2002). Moreover, the first *Chandra* observation of 1E 1613 in a low state hinted at a possible periodicity at ∼6 hours (Garmire et al. 2000) that was not confirmed by subsequent observations of the source in bright states. A long (90 ks) observation with *XMM-Newton*, performed in 2005, caught 1E 1613 in a low state and yielded unambiguous evidence for a strong, nearly sinusoidal modulation at $6.67 \pm 0.03$ hours ($24.0 \pm 0.1$ ks; De Luca et al. 2006). The same periodicity was then recognised also in the older data-sets, albeit with a very different pulse shape, including two narrow dips per period. No faster pulsations are seen in 1E 1613 (De Luca et al. 2006).

Large flux variations, similar to those observed in 1E 1613, are common among magnetars (e.g. Rea & Esposito 2011), but these pulsars, whose emission is believed to be powered mainly by the magnetic field, are characterised by rotational periods in the narrow range 2–12 s. On the other hand, CCOs are steady sources, and their periods –when known– are in the 0.1–0.5 s range (Zavlin et al. 2000; Gotthelf, Halpern & Seward 2005; Gotthelf & Halpern 2009). If 1E 1613 is indeed a magnetar, it must have been slowed down by some unusual mechanisms, perhaps by a propeller interaction with a debris disk (De Luca et al. 2006; Li 2007). A different possibility is that 1E 1613 is a peculiar low-mass binary,[1] powered by a double (wind plus disk) accretion onto a

---

⋆ E-mail: paoloesp@oa-cagliari.inaf.it

[1] Deep observations of the field of 1E 1613 with the Very Large Telescope and the *Hubble Space Telescope* showed only two or three faint infrared sources ($H \sim 22$) consistent with the position of 1E 1613. If none of them

recently-born compact object (in this case the 24-ks signal would result from the orbital motion of the system) or hosting a magnetar (De Luca et al. 2006; Pizzolato et al. 2008; Bhadkamkar & Ghosh 2009). Both scenarios require nonstandard assumptions about the formation and evolution of compact objects in supernova explosions.

Here we report on the results from a 5-year monitoring of 1E 1613 with the *Swift* satellite (Gehrels et al. 2004). This unique data set allowed us to obtain a phase-coherent timing solution encompassing also *Chandra* and *XMM-Newton* archival observations. Thanks to this, we are able to derive an accurate period for 1E 1613 and to set the first upper limits on the period derivative for this puzzling source.

## 2 X-RAY OBSERVATIONS AND DATA REDUCTION

### 2.1 *Swift* data

The X-Ray Telescope (XRT; Burrows et al. 2005) on-board *Swift* uses a front-illuminated CCD detector sensitive to photons between 0.2 and 10 keV with an effective area of about 110 cm$^2$ (at 1.5 keV) and a field view of 23-arcmin in diameter. Two main readout modes are available: photon counting (PC) and windowed timing (WT). PC mode provides two dimensional imaging information and a 2.5073-s time resolution; in WT mode only one-dimensional imaging is preserved, achieving a time resolution of 1.766 ms.

Between April 2006 and April 2011, 1E 1613 was observed by XRT 49 times, for a total net exposure time of 102.8 ks in PC mode.[2] The distribution of the *Swift* observations can be seen in the long-term light curve in Fig. 1, while in Fig. 2 we show the image of 1E 1613 and RCW 103 resulting from all the XRT data gathered so far. Except for periods in which the source was not visible by the XRT because of pointing constraints of the *Swift* spacecraft, approximately one 2-ks observation in imaging mode was collected per month. On a few occasions, when 1E 1613 showed hints of consistent flux variations, we requested target-of-opportunity observations. For example this happened at the end of October 2010 (see Fig. 1 around MJD 55 500), when the source count rate remained at a relatively high level of $\approx$0.09 counts s$^{-1}$ for a few consecutive pointings spanning $\sim$5 days. The observations were not time-constrained, so the monitoring can be considered a *casual* sampling of the phase of 1E 1613.

The XRT data were uniformly processed with XRTPIPELINE (version 12, in the HEASOFT software package version 6.9), filtered and screened with standard criteria. In order to reduce the contamination from the SNR, the source counts were energy-selected in the 1–10 keV band and extracted within a 10-pixel radius (one XRT pixel corresponds to about $2\farcs36$). To convert the photon arrival times to the Solar system barycentre for the timing analysis, we used the BARYCORR task and the *Chandra* position of De Luca et al. (2008). For the spectroscopy, we used the latest spectral redistribution matrices in CALDB (20091130), while the ancillary response files were generated with XRTMKARF, which accounts for different extraction regions, vignetting and point-spread function corrections.



**Figure 1.** *Swift*/XRT light curve of 1E 1613 (1–10 keV). Each observation is represented by a point in the plot. A rough conversion of 0.1 count s$^{-1}$ $\simeq$ $5 \times 10^{-12}$ erg cm$^{-2}$ s$^{-1}$ (1–10 keV, not corrected for absorption) can be derived from the spectral analysis (Section 3).



**Figure 2.** *Swift*/XRT image of 1E 1613 and its SNR, RCW 103. All the XRT data (as of 2011 April) were combined, totalling to 102.8 ks exposure time. Photon energy is colour-coded: red corresponds to 0.2–0.9 keV energies, green to 0.9–1.7 keV, blue to 1.7–8.0 keV. North is up, east is left. The source and background regions considered for the analysis (Section 3) are over-plotted.

### 2.2 *Chandra* and *XMM-Newton* data

We complemented the *Swift*/XRT data-set with the few *Chandra* and *XMM-Newton* observations long-enough to contain a minimum of two modulation cycles of 1E 1613 (Table 1). The *Chandra*/ACIS-S observation (performed on 2002 March 03, when the source was rather bright) has been already published in Sanwal et al. (2002), the *XMM-Newton*/EPIC (2005 August 23–24) and *Chandra*/HRC-S (2007 July 03) ones, both carried out while 1E 1613 was in a low state, have been published in De Luca et al. (2006) and in De Luca et al. (2008), respectively; we refer to these papers for more details. The *Chandra*/ACIS-I observation of 2010 June 01 is reported here for the first time. For this work the data were processed and analysed with standard procedures, using the latest available versions of the *Chandra* Interactive Analysis of Observation software (CIAO, version 4.2) and of the *XMM-Newton* Science Analysis Software (SAS, version 10).

---

is linked to the X-ray source, 1E 1613 is undetected in the near infrared down to $H > 23$ (De Luca et al. 2008).

[2] Also 5.5 ks of WT data were collected during the same pointings. However, given the bright SNR in which 1E 1613 is embedded (see Fig. 2), we did not make use of them in this work.

**Table 1.** *XMM-Newton* and *Chandra* observations used for this work.

| Instrument | Obs.ID | Date$^a$ (MJD TBD) | Duration$^b$ (ks) |
|---|---|---|---|
| *Chandra*/ACIS-S | 2759 | 52336.489 | 50.3 |
| *XMM-Newton*/EPIC | 0302390101 | 53605.824 | 87.5 |
| *Chandra*/HRC-S | 7619 | 54284.330 | 80.2 |
| *Chandra*/ACIS-I | 11823 | 55348.604 | 62.5 |

$^a$ Mid-point of observation.

$^b$ Time between first and last event.

## 3 ANALYSIS AND RESULTS

We merged the data from the *Swift*/XRT observations and accumulated a combined spectrum (a detailed spectral analysis will be reported elsewhere). The data were rebinned with a minimum of 20 counts per energy bin and the background was estimated from an annular region centred on 1E 1613 (with radii 10 and 20 pixel, see Fig. 2). The spectrum can be fit [$\chi^2_\nu = 0.82$ for 174 degrees of freedom (dof)] by a double-blackbody corrected for the interstellar absorption (see De Luca et al. 2006). The best-fit parameters are blackbody temperatures $kT_1 = 0.50^{+0.06}_{-0.11}$ keV and $kT_2 = 0.8^{+0.5}_{-0.1}$ keV, radii $R_1 = 0.6^{+0.2}_{-0.1}$ km and $R_2 = 0.12^{+0.10}_{-0.09}$ km (for a distance of 3.3 kpc; Caswell et al. 1975), and absorption $N_{\rm H} = (7^{+1}_{-2}) \times 10^{21}$ cm$^{-2}$ ($1\sigma$ errors). The observed averaged flux is $\sim$$1.7 \times 10^{-12}$ erg cm$^{-2}$ s$^{-1}$, similar to that measured with *XMM-Newton* in the August 2005 (De Luca et al. 2006).

As can be seen from Fig. 1, 1E 1613 showed only moderate variability between the many *Swift* pointings, remaining always well below the flux level of the 1999–2000 outburst ($\sim$$5 \times 10^{-11}$ erg cm$^{-2}$ s$^{-1}$; Garmire et al. 2000). Also the *Chandra* HRC-S and ACIS-I observations show a flux of $\approx$$2 \times 10^{-12}$ erg cm$^{-2}$ s$^{-1}$. We note that on a statistical ground we are sensitive only to outbursts lasting $\gtrsim$ 1 month. Although shorter phases of enhanced emission cannot be ruled out for 1E 1613 based on our data, the historical behaviour of the source indicates that its outbursts are likely longer than a few months (De Luca et al. 2006). Thus our data suggest that 1E 1613 persisted in a quiescent state during the five years of the *Swift* monitoring.

Owing to their long time span (from MJD 53 804.500 to 55 632.581), the *Swift* data are suitable for studying the 24-ks modulation of 1E 1613. A folded profile showing a significant modulation cannot be obtained from the XRT data using the most accurate period available so far, i.e. the one estimated from the August 2005 *XMM-Newton* observation ($P = 24.0 \pm 0.1$ ks; De Luca et al. 2006). This is not surprising since $\sim$200 days separate the *XMM-Newton* observation from the start of the *Swift* monitoring, and the *XMM-Newton* period uncertainty implies a phase uncertainty of half a cycle after only $\sim$30 days.

So, as a starting point, we computed a fast-Fourier-transform power spectrum using all the *Swift* data at the highest resolution allowed by the PC mode (bin time 2.5073 s). Given the approximate knowledge of the source period, a 'blind' search is not, in principle, necessary, but we did this to have a clear picture of the *Swift* time-series: considerable noise can be expected to result from both source flux variations and the *Swift* uneven sampling of the light curve, and a search restricted around the *XMM-Newton* period would have involved the risk of selecting a spurious signal, in the case the true signal was embedded in a high level of non-white noise. As expected, significant noise is present, but a very prominent peak at 24 031(2) s (the quoted uncertainty indicates the Fourier period resolution) stands out well above the noise level, with a Leahy-normalised power (Leahy et al. 1983) of 425 (Fig. 3). While the non-white noise does not affect the frequency of a real signal, it alters the statistical properties of a time-series; following the recipes of Israel & Stella (1996) we estimate that, after taking into account the number of frequencies searched, the probability of having a signal this strong by chance coincidence is lower than $3 \times 10^{-9}$ (that is a detection at a higher than 5.9$\sigma$ confidence level). Moreover, the signal is consistent with the periodicity measured by *XMM-Newton*. We also note that no other periodicity with significance higher than 3$\sigma$ was found up to $\sim$1 year.



**Figure 3.** Power spectrum computed for the combined *Swift*/XRT data. The peak corresponding to the 24-ks signal is evident.

We folded the *Swift* data, as well as those of the observations in Table 1, on the period $P = 24\,031$ s. We obtained very significant pulse profiles; those of the data taken during the quiescent state of 1E 1613 (*Swift*, *XMM-Newton*, *Chandra*/HRC-S and ACIS-I) are single-peaked (and well-modelled by two or three sine functions with the periods fixed at the fundamental period and higher harmonics, with phases and amplitudes free to vary), while that from the *Chandra*/ACIS-S observation shows two asymmetric peaks per cycle, both exhibiting two sub-peaks. In order to obtain a refined ephemeris, we studied the phase evolution through the epoch-folded data by means of an iterative phase-fitting technique (see e.g. Dall'Osso et al. 2003). Given the variability of the pulse shape, we did not make use of a pulse template to cross-correlate with, but we inferred the phase of the modulation by fitting each individual folded profile with the fundamental plus three higher harmonics.[3]

As a first step, we fit the phases of fundamental harmonic obtained from the *Swift* data divided into four segments of approximately equal length and from the *XMM-Newton* and *Chandra* HRC-S and ACIS-I observations. At this stage, the *Chandra*/ACIS-S observation was left out, because of its very different pulse profile. The time-evolution of the phase can be followed unambiguously throughout all the data and described with a linear relation of the form $\phi = \phi_0 + 2\pi(t - t_0)/P$. We assumed the start of the *Swift* monitoring, MJD 55 804.0, as the reference epoch $t_0$ and the fit ($\chi^2_\nu = 0.88$ for 5 dof) gives $P_{\rm A} = 24\,030.42(2)$ s (1$\sigma$ uncertainty, valid over the range MJD 53 605–55 632). We designate this rotational ephemeris 'solution A'. A quadratic term $-\pi(t - t_0)^2 \dot{P}/P^2$, which would reflect the presence of a period derivative ($\dot{P}$), is

---

[3] This is the minimum number of harmonics necessary to properly fit all the folded profiles, including the multi-peaked *Chandra*/ACIS-S one. The timing analysis described in the following was performed also with a slightly different approach: the phases of the individual folded profiles were derived by a fit with a variable number of harmonics, determined on a case-by-case basis by requesting that the addition of a further (higher) harmonic is not statistically significant (by means of a Fisher-test) with respect to the null hypothesis; the results are essentially identical.



**Figure 4.** 32-bin epoch-folded pulse profiles of 1E 1613 obtained from different instruments (as indicated in each panel). The blue line distinguishes the *Chandra*/ACIS-S data, which were used in deriving solution B but not for solution A (see Section 3 for details). In red we over-plotted the *XMM-Newton*/EPIC data before the filtering for proton flares. The fundamental harmonic of the pulse profile is shown in green in each panel (in the EPIC panel it refers to the filtered data).

not required. This implies an upper limit on the period derivative of 1E 1613 of $|\dot{P}_{\rm A}| < 3.3 \times 10^{-9}$ s s$^{-1}$ ($3\sigma$ confidence level). In Fig. 4 we show the epoch-folded pulse profiles (including the *Chandra*/ACIS-S one) obtained using this solution. We note that the *XMM-Newton* observation was affected by rather intense proton flares. The removal of the intervals of flaring background, because of the few cycles contained in the observation, significantly affects the pulse profiles. As a check of the robustness of our results, we repeated the timing analysis using the unfiltered EPIC data and we obtained virtually identical results (both filtered and unfiltered profiles are plotted in Fig. 4).

Using solution A, we are able to predict for 1E 1613 the phase of the fundamental harmonic at the epoch of the *Chandra*/ACIS-S observation within $\pm 0.03$ cycles (at $3\sigma$). The phase of the fundamental harmonic *measured* in the Chandra/ACIS-S data nicely dovetails with the predicted value. Thus we derived a new coherent timing solution, which we denote with ‘B’, including the *Chandra*/ACIS-S data and therefore valid over the range MJD 52 336–55 632. Again, the phases of the fundamental harmonic can be fit with a linear relation ($\chi^2_\nu = 0.77$ for 6 dof) which yields $P_{\rm B} = 24\,030.42(2)$ s ($1\sigma$ uncertainty; epoch MJD 55 804.0). While the best-fitting period is equal to that of solution A, the limit on the period derivative is slightly more constraining: $|\dot{P}_{\rm B}| < 1.6 \times 10^{-9}$ s s$^{-1}$ ($3\sigma$ confidence level).

## 4 DISCUSSION

We presented the analysis of the first five years (2006 April–2011 April) of the *Swift*/XRT monitoring of the enigmatic X-ray source 1E 1613 at the centre of the SNR RCW 103. During this time span, the source remained in a quiescent state, with an average observed 1–10 keV flux of $\sim 1.7 \times 10^{-12}$ erg cm$^{-2}$ s$^{-1}$, 20–30 times lower than the historical maximum attained in the 1999–2000 outburst ($\sim 5 \times 10^{-11}$ erg cm$^{-2}$ s$^{-1}$; Garmire et al. 2000). The timing study of the *Swift* data yielded an accurate measure of the modulation period of 1E 1613 which allowed us to phase-connect the XRT data with archival *Chandra* and *XMM-Newton* observations. We derived two timing solutions, labelled A and B, both consistent with the same constant period $P_{\rm A} = P_{\rm B} = 24\,030.42(2)$ s but with somewhat different upper limits on the period derivative. Solution A ($|\dot{P}_{\rm A}| < 3.3 \times 10^{-9}$ s s$^{-1}$; MJD 53 605–55 632) is based on the single-peaked pulse profiles observed while 1E 1613 was in quiescence. Solution B ($|\dot{P}_{\rm B}| < 1.6 \times 10^{-9}$ s s$^{-1}$; MJD 52 336–55 632) is a natural extension of solution A including the multi-peaked *Chandra*/ACIS-S profile obtained on 2002 March 03, when 1E 1613 was in a rather bright state (Sanwal et al. 2002).

The phase-coherent timing technique employed in our timing study of 1E 1613 closely parallels the well-tested (virtually all systematics are under control) procedures used for X-ray bright spin-powered pulsars. For such objects it is implicitly assumed that the pulse profile does not intrinsically change in time, so that, when a Fourier decomposition of the pulse is introduced, neither the fundamental nor the higher harmonics evolve. In such a picture all the variability is due to Poisson noise and the phase evolution can be tracked by matching the pulse profile at any given epoch always with the same template.

In the case of 1E 1613 there are clear indications that the pulse shape changes in time so that the previous assumptions are not valid. This forced us to abandon the standard template-matching analysis and follow instead the phase of the fundamental (first) harmonic.[4] The main justification for such an approach is that usually (moderate) pulse shape changes are associated with higher harmonics while the fundamental is stable. Even if this seems to be the case for 1E 1613, given the stability of the phase of the fundamental harmonic over several years, we stress that it does not have to hold in general (see, e.g., Hartman et al. 2008).

In the following we discuss the implications of our newly derived upper limit $|\dot{P}| < 1.6 \times 10^{-9}$ s s$^{-1}$ on the models that have been proposed so far for 1E 1613. We remark that all the ensuing considerations are based on the assumption (discussed above) that the fundamental harmonic is a good tracer of the timing behaviour of the source.

Although the nature of 1E 1613 is still an open issue, most interpretations favour the neutron star scenario, in which the star is either isolated (De Luca et al. 2006; Li 2007) or in a binary system with a low-mass companion (Pizzolato et al. 2008; Bhadkamkar & Ghosh 2009). While the model by Bhadkamkar & Ghosh (2009) is based on a fast-spinning, moderately magnetised neutron star in a 6.67 hr eccentric orbit, in all the other cases an ultra-magnetised neutrons star ($B \approx 10^{15}$ G) is required to explain the very long period of 1E 1613 and the observed X-ray periodicity is related to the star spin period (in the binary scenario of Pizzolato et al. 2008 the latter may or may not coincide with the orbital period). Actually, as already noted by De Luca et al. (2006), magneto-dipolar braking alone is not enough to explain the present value of $P$ even invoking a magnetar, and spin-down by the interaction of the star magnetosphere with a (residual) disc is also necessary for an isolated object. Spin-down to $P \simeq 6.67$ hr in a time ∼2 kyr can be achieved if the initial period is peculiarly long ($\gtrsim$300 ms, De Luca et al. 2006; magnetars are in fact believed to be born with ms periods, Thompson & Duncan 1993). Using different assumptions on the termination radius of the disc (see Ekşi & Alpar 2005), Li (2007) showed that the same result can be recovered also for more conventional values of the initial period (≈10 ms).

---

[4] The large phase uncertainties do not allow a similar study of the higher harmonics (which are not even always detectable).

The current spin-down rate expected in the binary magnetar model is very small since the equilibrium period is reached well in advance of ∼1000 yr, unless the synchronisation time is very short (∼10 yr) and there is no mass transfer in the system (Pizzolato et al. 2008). If 1E 1613 is an object of this kind our upper limit on $\dot{P}$ is completely non-constraining for the model.

On the other hand, if 1E 1613 is an isolated magnetar surrounded by a fossil disc it must be a quite rare system. According to the Monte Carlo simulations of Li (2007), the fraction of objects of this type with periods $\gtrsim$100 s at an age of 2.5 kyr is only ∼1%. The large majority of stars are much faster rotators ($P \sim$ 1–10 s) in the ejector stage (typically identified with SGRs/AXPs) while ultra-slow systems are in the propeller/accretor phase.[5]

The spin-down rate of a neutron star which interacts with a fossil disc in the propeller stage is given by $\dot{P} \sim -\dot{M}R_{in}^2[\Omega_K(R_{in}) - 2\pi/P]P^2/(\pi I)$ where $R_{in} = 0.5[B^4R^{12}/(8GM\dot{M}^2)]^{1/7}$ is the Alfvén radius, $M \sim 1.4M_{\odot}$ and $R \sim 10^6$ cm are the star mass and radius ($I \sim 10^{45}$ g cm$^2$ is the moment of inertia), $B$ is the surface magnetic field, $\Omega_K$ is the Keplerian angular velocity and $\dot{M} \propto t^{-1.25}$ is the mass loss rate from the disc (Li 2007, and references therein).[6] If 1E 1613 is currently in the propeller phase, $\dot{P}$ can be derived from the previous expressions with $P \sim 24$ ks and $t \sim 2.5$ kyr, as a function of the disc initial mass $M_D(0)$ and of $B$. Results are shown in Fig. 5 (lower panel) for $10^{-6} < M_D(0)/M_{\odot} < 10^{-2}$ and $3.2 \times 10^{13}$ G $< B < 10^{16}$ G; the Alfvén radius as a function of $M_D(0)$ and $B$ is plotted in the upper panel. Comparison of $R_{in}$ with the light cylinder radius, $R_{LC} = cP/(2\pi)$, and the co-rotation radius, $R_C = [GMP^2/(4\pi^2)]^{1/3}$ shows that the star is currently in the propeller phase, characterised by $R_C < R_{in} < R_{LC}$, only if $B \gtrsim 10^{15}$ G. Lower values of the field or large enough (depending on $B$) initial disc mass would result in 1E 1613 being in the accretor stage. The resulting values of $\dot{P}$ are orders of magnitude above the upper limit we reported (as a reference, the case discussed in De Luca et al. 2006 has $\dot{P} \sim 10^{-7}$ s s$^{-1}$), with the only exception of an extremely narrow range of $M_D(0)$ for each $B$. Although not all the values of $M_D(0)$ and $B$ we considered will produce the correct period at an age of 2.5 kyr (this depends also on both the initial period and the angle between the magnetic and spin axes, see again Li 2007), our conclusion is that if 1E 1613 is an isolated magnetar, and hence it must be either a propeller or an accretor at present, the typical large expected values of $\dot{P} \gg 3 \times 10^{-9}$ s s $^{-1}$ makes the propeller option rather unlikely. Assessing the reliability of the accretor scenario requires a more thorough analysis. Here we just note that if the same expression of the torque remains valid during the accretion phase, the spin-up rate is also largely in excess of our upper limit on $\dot{P}$.



**Figure 5.** Upper panel: the Alfvén radius as a function of the initial disc mass $M_D(0)$ for different values of the magnetic field in the range $13.5 \leqslant \log B\,(\mathrm{G}) \leqslant 16$ (the field increases as the curve moves upwards); the light cylinder radius $R_{LC}$ and the corotation radius $R_C$ are also shown. Lower panel: the period derivative as a function of $M_D(0)$ for those values of $B$ for which the source is currently in the propeller stage; the dashed line marks our derived upper limit on $\dot{P}$. Here the period and age have been fixed to those inferred for 1E 1613 (see text).

## ACKNOWLEDGMENTS

We thank the referee, David Helfand, for his constructive and valuable comments. This research is based on data and software provided by the NASA/GSFC's High Energy Astrophysics Science Archive Research Center (HEASARC), the *Chandra* X-ray Center (CXC, operated for NASA by SAO), and the ESA's *XMM-Newton*

---

[5] Although previous figures are model-dependent the conclusion that propellers/accretors are a tiny minority appears to be robust.

[6] We remark that no complete theory of the interaction of a disc with a magnetised neutron star exists. Following Li (2007) we adopted the 'efficient' form of the propeller torque (see e.g. Francischelli & Wijers 2002, for a comparison of different expressions of the torques).

Science Archive (XSA). PE acknowledges financial support from the Autonomous Region of Sardinia through a research grant under the program PO Sardegna FSE 2007–2013, L.R. 7/2007 “Promoting scientific research and innovation technology in Sardinia”. This work was partially supported by the ASI/INAF contract I/009/10/0.

## REFERENCES

Becker W., Aschenbach B., 2002, in Becker W., Lesch H., & Trümper J., eds., Proc. of the 270 WE-Heraeus Seminar, Neutron Stars, Pulsars, and Supernova Remnants Vol. 278 of MPE Report, Garching, p. 64
Bhadkamkar H., Ghosh P., 2009, A&A, 506, 1297
Burrows D. N., et al., 2005, Space Science Reviews, 120, 165
Carter L. M., Dickel J. R., Bomans D. J., 1997, PASP, 109, 990
Caswell J. L., Murray J. D., Roger R. S., Cole D. J., Cooke D. J., 1975, A&A, 45, 239
Dall’Osso S., Israel G. L., Stella L., Possenti A., Perozzi E., 2003, ApJ, 599, 485
De Luca A., 2008, in Bassa C., Wang Z., Cumming A., Kaspi V. M., eds., 40 years of pulsars: Millisecond Pulsars, Magnetars and More. Vol. 983 of AIP Conf. Proc., Melville NY, p. 311
De Luca A., Caraveo P. A., Mereghetti S., Tiengo A., Bignami G. F., 2006, Science, 313, 814
De Luca A., Mignani R. P., Zaggia S., Beccari G., Mereghetti S., Caraveo P. A., Bignami G. F., 2008, ApJ, 682, 1185
Ekşi K.Y., Alpar M.A. 2005, ApJ, 620, 390
Francischelli G. J., Wijers R. A. M. J. 2002, preprint (astro-ph/0205212)
Garmire G. P., Garmire A. B., Pavlov G., Burrows D. N., 2000, in AAS/HEAD Fifth Meeting, Vol. 32 of Bulletin of the American Astronomical Society, Washington DC, p. 1237
Garmire G. P., Pavlov G. G., Garmire A. B., Zavlin V. E., 2000, IAU Circ., 7350, 2
Gehrels N., et al., 2004, ApJ, 611, 1005
Gotthelf E. V., Halpern J. P., 2009, ApJ, 695, L35
Gotthelf E. V., Halpern J. P., Seward F. D., 2005, ApJ, 627, 390
Gotthelf E. V., Petre R., Hwang U., 1997, ApJ, 487, L175
Gotthelf E. V., Petre R., Vasisht G., 1999, ApJ, 514, L107
Hartman J. M., et al., 2008, ApJ, 675, 1468
Israel G. L., Stella L., 1996, ApJ, 468, 369
Leahy J. A., Darbo W., Elsner R. F., Weisskopf M. C., Sutherland P. G., Kahn S., Grindlay J. E., 1983, ApJ, 266, 160
Li X.-D., 2007, ApJ, 666, L81
Pizzolato F., Colpi M., De Luca A., Mereghetti S., Tiengo A., 2008, ApJ, 681, 530
Rea N., Esposito P., 2011, in Torres D. F., Rea N., eds, Astrophysics and Space Science Proceedings, High-Energy Emission from Pulsars and their Systems. Springer Berlin, p. 247
Sanwal D., Garmire G. P., Garmire A., Pavlov G. G., Mignani R., 2002, in 200th AAS Meeting, Vol. 34 of Bulletin of the American Astronomical Society, Washington DC, p. 764
Thompson C., Duncan R. C., 1993, ApJ, 408, 194
Tuohy I., Garmire G., 1980, ApJ, 239, L107
Tuohy I. R., Dopita M. A., Garmire G. P., Manchester R. N., 1983, ApJ, 268, 778
Zavlin V. E., Pavlov G. G., Sanwal D., Trümper J., 2000, ApJ, 540, L25

This paper has been typeset from a TEX/ LATEX file prepared by the author.